

Determination of antibiotics by SPE-LC-MS/MS in wastewater and risk assessment

Senar Aydın¹, Mehmet E. Aydın², Arzu Ulvi^{*1} and Havva Kılıç¹

¹Department of Environmental Engineering, Faculty of Engineering and Architecture,
Necmettin Erbakan University, 42140 Konya-Turkey

²Department of Civil Engineering, Faculty of Engineering and Architecture,
Necmettin Erbakan University, 42140 Konya-Turkey

(Received June 1, 2018, Revised February 8, 2019, Accepted February 20, 2019)

Abstract. In this study, conditions of solid phase extraction (SPE) for determination of some antibiotics such as trimethoprim, oxytetracycline, erythromycin, clarithromycin, azythromycin, doxycycline, sulfamethazine, ciprofloxacin, chlortetracycline, sulfamethoxazole in wastewaters were optimized. After the optimum volume and pH of the sample were determined, the effect of the concentration of the compounds and matrix were investigated. The highest recovery rates for antibiotic compounds were determined between 82% and 105% in 200 mL sample volume and pH 2.5. Then, antibiotic compounds were investigated in influent and effluent samples taken from Konya Urban Wastewater Treatment Plant. The concentration of the antibiotics was detected range of 0.11-101 ng/L in influent waters and <dl-288 ng/L in effluent samples in wastewater treatment plant. Hazard quotients (HQs) of antibiotic compounds determined in WWTP effluents to evaluate the risk towards different aquatic organisms (algae, *Daphnia magna* and fish) were determined. Azythromycin for fish and erythromycin, sulfamethoxazole, ciprofloxacin, clarithromycin for algae posed a moderate risk while azythromycin, ciprofloxacin, clarithromycin, oxytetracycline posed a high risk for algae in the receiving environment.

Keywords: antibiotic; LC-MS/MS; risk assessment; SPE, wastewater

1. Introduction

Pharmaceuticals are groups of chemical substance having medical features. Wastewater treatment plant effluents, groundwater and surface water may involve 160 types of drugs. In fact, even pharmaceuticals in drinking water are encountered. Nowadays, pharmaceuticals have become indispensable and important element in areas such as medicine, veterinary, agriculture. In the 1970s, information on the presence of pharmaceuticals in nature has begun to emerge in recent years and their effects on behavior studies were started in the 1990s. The main source of pharmaceutical compounds is the metabolic wastes of patients going to the sewerage system. These contaminants are excreted by fecal and urinary because of not to be fully metabolized by humans and animals. In generally, it is assumed that the hospital wastewater has the same pollutant

*Corresponding author, Ph.D., E-mail: atekinay@konya.edu.tr

- determination of 13 quinolone antibiotic derivatives in wastewater samples using solid phase extraction and ultraperformance liquid chromatography-tandem mass spectrometry”, *Microchem. J.*, **106**, 323-333.
- EPA Method 1694 (2007), *Pharmaceuticals and Personal Care Products in Water, Soil, Sediment, and Biosolids by HPLC/MS/MS*, December 2007.
- Gros, M., Rodríguez-Mozaz, S. and Barceló, D. (2013), “Rapid analysis of multiclass antibiotic residues and some of their metabolites in hospital, urban wastewater and river water by ultra-high-performance liquid chromatography coupled to quadrupole-linear ion trap tandem mass spectrometry”, *J. Chromatogr. A*, **1292**, 173-188.
- Guerra, P., Kim, M., Shah, A., Alaei, M. and Smyth, S.A. (2014), “Occurrence and fate of antibiotic, analgesic/anti-inflammatory, and antifungal compounds in five wastewater treatment processes”, *Sci. Total Environ.*, **473**, 235-243.
- Hirsch, R., Ternes, T., Haberer, K. and Kratz, K.L. (1999), “Occurrence of antibiotics in the aquatic environment”, *Sci. Total Environ.*, **225**(1-2), 109-118.
- Huerta, B., Rodríguez-Mozaz, S. and Barceló, D. (2012), “Pharmaceuticals in biota in the aquatic environment: Analytical methods and environmental implications”, *Anal. Bioanal. Chem.*, **404**(9), 2611-2624.
- Khan, G.A., Lindberg, R., Grabic, R. and Fick, J. (2012), “The development and application of a system for simultaneously determining anti-infectives and nasal decongestants using on-line solid-phase extraction and liquid chromatography-tandem mass spectrometry”, *J. Pharmaceut. Biomed. Anal.*, **66**, 24-32.
- Kümmerer, K. (2008), *Pharmaceuticals in the Environment; Sources, Fate, Effects and Risks*, Springer.
- López-Serna, R., Petrović, M. and Barceló, D. (2011), “Development of a fast instrumental method for the analysis of pharmaceuticals in environmental and wastewaters based on ultrahigh performance liquid chromatography (UHPLC)-tandem mass spectrometry (MS/MS)”, *Chemosphere*, **85**(8), 1390-1399.
- Petrovic, M., Gros, M. and Barcelo, D. (2006), “Multi-residue analysis of pharmaceuticals in wastewater by ultra-performance liquid chromatography-quadrupole-time of-flight mass spectrometry”, *J. Chromatogr. A*, **1124**(1-2), 68-81.
- Ramirez, A.J., Brain, R.A., Usenko, S., Mottaleb, M.A., O’Donnell, J.G., Stahl, L.L., Wathen, J.B., Snyder, B.D., Pitt, J.L., Perez-Hurtado, P., Dobbins, L.L., Brooks, B.W. and Chambliss, C.K. (2009), “Occurrence of pharmaceuticals and personal care products in fish: results of a national pilot study in the United States”, *Environ. Toxicol. Chem.*, **28**(12), 2587-2597.
- Tong, L., Liu, H., Xie, C. and Li, M. (2016), “Quantitative analysis of antibiotics in aquifer sediments by liquid chromatography coupled to high resolution mass spectrometry”, *J. Chromatogr. A*, **1452**, 58-66.
- Verlicchi, P., Al Aukidy, M., Galletti, A., Petrovic, M. and Barceló, D. (2012), “Hospital effluent: Investigation of the concentrations and distribution of pharmaceuticals and environmental risk assessment”, *Sci. Total Environ.*, **430**, 109-118.
- Yuan, X., Qiang, Z., Ben, W., Zhu, B. and Liu, J. (2014), “Rapid detection of multiple class pharmaceuticals in both municipal wastewater and sludge with ultrahigh performance liquid chromatography tandem mass spectrometry”, *J. Environ. Sci.*, **26**(9), 1949-1959.